

Figure 1






Duration of staurosporine treatment (h)



ANT TO COLOR




## Figure 1 (con't)



ESTERIS PETERS (1)








Figure 3





Figure 4



mt = K<u>ISVYLTAAVVGFVAYGIL</u>KWYRGT

dana ang mga Taneg 30 mg;


101904WCsq7070R

## Figure 5

					domain is a mitochondrial targeting signal	domain is required for function
M11L	K	ISVYLTAAVVGFVAYGIL	K	WYRGT	Y	Y
Bc1-2	1X	TLLSLALVGACITLGAYLS	H	K	Y	Y/N
$\mathtt{Bcl-X_L}$	R	WFLTGMTVAGVVLLGSLFS	R	K	Y	Y/N
Boo/Diva	R	LLIQAFLSGFFATAIFFIW	X.	RL	?	?
CED-9	R	WSMIGAGVTAGAIGIVGVVVCG	R	MMFSLE	, <b>5</b> ,,	
BHRF-1	ĸ	FSWTLFLAGLTLSLLVICSYLFI	s R	GRH	Y	Y
KSbcl-2	R	MTALLGSIALLATILAAVAMS	R	R	?	?
Nip3	K	VFLPSLLLSHLLAIGLGIYIG	R	RLTTSTSTI	· Y	Y
Nix	K	VFIPSLFLSHVLALGLGIYIG	X	RLSTPSA	Υ	Υ
	positive charge	18-24 aa putative membrane-spanning domain	positi charç			


进产用4人作为产品和2 等水溶解 30人人以3

Figure 6




TO BEAT PERMITTED

Figure 6 (con't)




GFP construct expressed by cells

Figure 7




CARLA TO A 100


Process to Identify M11L-interacting Proteins














Bak, a Bcl2 family member, was identified by Mass Spec

MASGAGPGPRAECGEPALPSASEEAVAAD
TEEVFRSYVFYRHQAEQEAEGVAAPADPEM
VTLPLAPSSTMGQVGRALAIIGDDINRRYD
SEFQTMLAHLAPTAENAYEYFTKIATSLFE
SGINWGRVVALLGFGYRLALHVYQH
GLTGFLGQVTRFVVDFMLHHCIARWIAQRGGWVA

ALNLGNGPILNVLVVLGVVLLGQFVVRRFFKS

Carlot Al Section 10

Figure 11